

Table 14. Reports of antiretroviral medications, by selected characteristics of persons with HIV infection (N = 1,622), Supplement to HIV/AIDS Surveillance, June 1997–December 2000

	Antiretroviral medications					
	Never		Taken in the past, but not currently		Currently taking	
	Number	%	Number	%	Number	%
Race/ethnicity						
White, not Hispanic						
Men	108	32.0	32	9.5	197	58.5
Women	31	33.0	10	10.6	53	56.4
Black, not Hispanic						
Men	199	39.0	38	7.5	273	53.5
Women	140	36.4	30	7.8	215	55.8
Hispanic						
Men	49	34.0	9	6.3	86	59.7
Women	18	19.4	12	12.9	63	67.7
Age (years)						
18–29	154	39.1	39	9.9	201	51.0
30–39	214	33.1	73	11.3	359	55.6
40–49	144	33.0	24	5.5	268	61.5
≥50	54	37.0	3	2.0	89	61.0
Exposure category						
Men						
Male-to-male sexual contact	187	32.9	49	8.6	332	58.5
Injection drug use	39	31.0	12	9.5	75	59.5
Male-to-male sexual contact with injection drug use	49	41.5	14	11.9	55	46.6
Heterosexual contact	59	47.6	7	5.6	58	46.8
Women						
Injection drug use	46	37.4	12	9.8	65	52.8
Heterosexual contact	110	34.3	33	10.3	178	55.4
Most recent CD4 count						
<200	10	16.7	12	20.0	38	63.3
200–499	107	25.0	36	8.4	285	66.6
≥500	196	33.2	53	9.0	342	57.8
Don't know/never had CD4 count	187	40.3	36	7.8	241	51.9
Interview period						
1997–1998	295	40.9	68	9.4	359	49.7
1999–2000	271	30.1	71	7.9	558	62.0
Overall	566	34.9	139	8.6	917	56.5

Note. Excludes persons for whom data were missing. Values may not sum to totals because missing data.